

Prenatal and Postnatal Care

Access to Prenatal Care in Rural Northern California

- Healthy pregnancies occur when comprehensive, routine prenatal care begins early in pregnancy. Receiving prenatal care during the first trimester improves maternal and infant health. Women who do not receive prenatal care are at almost three times the risk of having a low-birthweight infant and increasing risk for poorer health outcomes.
- Women who are uninsured or those with no regular source of care prior to pregnancy are more likely to enter into prenatal care after their first trimester.^{1,2}
- Smoking and alcohol use in the three months before pregnancy and during pregnancy are more prevalent health behaviors of women in rural Northern California than in other areas of the state.³
- One of the most common complications for pregnant and postpartum women is depression. Studies show 14-23% of pregnant woman and as much as 25% of postpartum women experience depression.⁴

How Health Centers Provide the Necessary Care

Clinical Interventions

- Offer contraceptive services, pregnancy testing, and preconception counseling for all reproductive age women.
- Recommend that all reproductive age women take a multi-vitamin that includes a folic acid supplement. Adequate intake of folic acid may help prevent some birth defects.
- Ask all pregnant patients about tobacco, alcohol and other drug use and provide appropriate counseling or treatment interventions.
- For Partnership members, introduction to PHC's Maternal Child Health Program/Growing Together Program (GTP), which offers incentives for timely prenatal and postpartum care.
- Schedule postpartum visit 3-5 weeks after delivery to allow time for rescheduling if needed.
- Piggyback scheduling of infant and postpartum visits (if same provider) during prenatal visits and prior to hospital discharge.

Community Interventions

- Share patient handouts, brochures, or videos at community health fairs to raise awareness about everyday strategies to support healthy pregnancy, such as eating a balanced diet, staying active, and taking nutritional supplements.
- Provide health insurance information, public benefit programs, and enabling service enrollment support to patients.

¹ Kaiser Family Foundation. Promoting Access to Prenatal Care: Lessons from the California Experience. Spring 2003.

² Braveman P et al., Barriers to timely prenatal care among women with insurance: the importance of pregnancy factors, *Obstetrics & Gynecology*, 2000, 95(6):874-880.

³ Ibid.


⁴ Kaiser Family Foundation. Medicaid and Health Coverage for Low-Income Women in Pregnancy and After Childbirth. 10 Sep 2019.

Prenatal and Postnatal Care

Rural Northern California Health Center Data

Key Points

- Health centers often have pregnant women come in for prenatal care as new patients; some don't seek care until after their first trimester.
- Women choosing alternative care from a midwife or other practitioner may come to the health center during pregnancy for screenings or medical attention their primary practitioner cannot offer.
- Many Health centers in rural communities provide prenatal care to a relatively small population of women. Even one late entry can have a significant impact on these results.



Quality Measure Definitions (UDS)

The percentage of prenatal care patients who entered treatment during their first trimester.

- The Institute of Medicine estimates that every \$1 invested into proper prenatal care results in a savings of \$3.37 in neonatal care.⁵
- Maternal hospital stays with pregnancy and delivery-related complicating conditions account for \$17.4 billion in hospital costs in the U.S.⁶

National and State Quality Benchmarks

UDS 2021 CA Average: The average performance among health centers in California was 77.1%.

UDS 2021 U.S. Average: The average performance among health centers across the U.S. was 74.1%.

⁵ Lantos JD, Lauderdale DS. What is Behind the Rising Rates of Preterm Birth in the United States? RMMJ

⁶ Elixhauser A. (AHRQ) and Wier LM. (Thomson Reuters). *Complicating Conditions of Pregnancy and Childbirth, 2008*. HCUP Statistical Brief #113. May 2011. Agency for Healthcare Research and Quality, Rockville, MD. <http://www.hcup.us.ahrq.gov/reports/statbriefs/sb113.pdf>